

# Popantel<sup>®</sup> F Allwormer for Dogs

## Presentations:

10 kg dog tablets: - 2, 4 and 6 tablet display cards and 100 tablet bulk cartons  
 35 kg dog tablets: - 2 tablet display cards and 70 tablet bulk cartons

## Active Constituents:

FEBANTEL, PYRANTEL and PRAZIQUANTEL

(10 kg dose – 250 mg, 49.8 mg and 50 mg respectively; 35 kg dose – 875 mg, 174.4 mg and 175 mg respectively)

## Actions:

Popantel F is a treatment for gastro-intestinal parasites of dogs. Popantel F is effective against susceptible intestinal helminths and giardia. Popantel F is not a preventative treatment for canine heartworm disease.

## Indications:

For the control of ROUNDWORMS (*Toxocara canis*, *Toxascaris leonina*), WHIPWORMS (*Trichuris vulpis*) HOOKWORMS (*Ancylostoma caninum*, *A. braziliense*, *Ancylostoma ceylanicum*, *Uncinaria stenocephala*. TAPEWORMS (*Dipylidium caninum*, *Taenia ovis*, *T. pisiformis*, *T. hydatigena* and *Echinococcus granulosus* (hydatid tapeworm)). Also controls the shedding of oocysts of the protozoan *Giardia* sp. Does not control heartworm.

## Dosage and Administration:

POPANTEL<sup>®</sup> F is given as a single dose by mouth. Fasting prior to dosing is NOT necessary. IT IS ESSENTIAL TO WEIGH DOGS BEFORE TREATMENT. **If worm problems persist, consult a veterinarian.**

### Dosage:-

#### 10 kg Popantel F Tablets

Bodyweight	Tablets
1.5 – 2.5 kg	¼
2.6 – 5 kg	½
5.1 – 10 kg	1
Over 10 kg	1 tablet for every 10kg bodyweight. Round off doses to the nearest ½ tablet.

#### 35 kg Popantel F Tablets

Bodyweight	Tablets
20 – 35 kg	1
35.1 – 52 kg	1½
52.1 – 70 kg	2
70.1 – 87 kg	2½
Over 87 kg	3

**HOOKWORMS & ROUNDWORMS**

Dogs and puppies: treat at 2, 4, 8 and 12 weeks and then every month until six months of age; thereafter every 3 months or on veterinary advice. Pregnant Bitches: treat prior to mating, then every 10 days before whelping and at 2 and 4 weeks after whelping.

**WHIPWORMS**

Treat every 6 to 8 weeks after 3 months of age.

**TAPEWORMS** (excluding hydatid tapeworms)

Treat every 3 months.

**HYDATID TAPEWORM**

Dogs should not be fed, or allowed to feed on, offal from any species. Dogs in hydatid areas should be treated every 6 weeks. It should be noted that the hydatid tapeworm poses a severe risk to human health by transmission to humans from infected dogs.

**GIARDIA**

Suspected infection should be confirmed by a veterinarian. Treat for three consecutive days. Repeat treatments may be necessary on veterinary advice.

**First Aid:**

If poisoning occurs, contact a doctor or Poisons Information Centre. *Phone Australia 131126.*

Additional information is listed in the Material Safety Data Sheet available from Jurox Pty Ltd in Australia (1800 023 312)

**Scheduling:**

Not Scheduled

For Animal Treatment Only