



TECHNICAL NOTES

Ropel

Liquid Testosterone



Active Constituents

Testosterone enanthate 75 mg/mL (equivalent to 54 mg/mL Testosterone)

Indications

Prevention and treatment of external preputial ulceration and sheath rot in wethers.

Precautions

Avoid carcass damage.

Sterilise all injection apparatus by boiling before use. Avoid use of strong disinfectants on apparatus.

Avoid as far as possible, injection of animals during wet weather or under dusty conditions. The product must be injected only under the skin using the shortest needle possible, not exceeding 15 mm.

Dosage and Administration

For prevention:

Inject 1 mL subcutaneously high on the neck behind the ear.

Note: Prevention of external preputial ulceration and sheath rot is achieved by administering Ropel Liquid prior to the onset of the spring or autumn flush where wethers are grazing high nitrogen pastures.

For treatment:

Inject 2 mL subcutaneously high on the neck behind the ear. Repeat every 21 days as required.

Note: Where ulceration or sheath rot has developed, use the treatment dose of 2 mL injected subcutaneously and move the animals to other low nitrogen pastures where possible.





TECHNICAL NOTES



Withholding Periods

Meat: DO NOT ADMINISTER less than 21 days before slaughter for human consumption.

Trade Advice

Export Slaughter Interval (ESI):

This product does not have an ESI established.

For advice on the ESI, contact Jurox Pty Limited on 1800 023 312 before using this product.

First Aid

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 13 1126.

Presentation

200 mL pillow pack

Storage

Store below 25°C (air conditioning). Protect from light.

Poisons Schedule

S4

Registration Number

APVMA No. 38228

AU - Tech Note - Ropel / Nov 2018 / APPROVED



www.jurox.com.au
Customer Service 1800 023 312
® Registered Trademark of Jurox Pty Limited

AUSTRALIA'S
ANIMAL HEALTH COMPANY

